



---

**Environmental Health & Safety**

1314 Kinnear Rd.  
Columbus, Ohio 43212  
Phone (614) 292-1284  
Fax (614) 292-6404  
<http://www.ehs.osu.edu/>

## **Heat and Cold Stress Program**

Prepared by:  
The Ohio State University  
Environmental Health and Safety  
Occupational Safety & Industrial Hygiene

1314 Kinnear Road  
Columbus, OH 43212-1168

614-292-1284 Phone  
614-292-6404 Fax

[www.ehs.osu.edu](http://www.ehs.osu.edu)

**Table of Contents**

**1.0 Introduction .....3**

**2.0 Responsibilities.....3**

**3.0 Heat Related Illnesses; Signs, Treatment & Prevention .....4**

**4.0 Cold Related Illnesses; Signs, Treatment & Prevention .....6**

**5.0 Training .....9**

**6.0 Recordkeeping .....9**

**Appendix A: Heat Index.....10**

**Appendix B: Cold Stress Equation .....14**

## 1.0 Introduction

- 1.1 Working in extreme temperatures (hot or cold) can overwhelm the body's internal temperature control system. When the body is unable to warm or cool itself, heat or cold related stress can result. Heat and cold stress can contribute to adverse health effects which range in severity from discomfort to death.
- 1.2 Environmental Health and Safety (EHS) has developed this Heat and Cold Stress Program to minimize the effects of heat and cold stress on Ohio State University (OSU) employees. This program contains the procedures and practices for safely working in temperature extremes.
- 1.3 The Occupational Safety and Health Administration (OSHA) does not currently have specific standards for heat or cold stress. However, the Occupational Safety and Health Act of 1970 General Duty Clause (Section 5(a)(1)) states that *"Each employer shall furnish to each of his employees employment and a place of employment which are free from recognized hazards that are causing or are likely to cause death or serious physical harm to his employees."* In addition, 29 CFR Subpart I relating to personal protective equipment requires employers to provide protection to employees exposed to hazards in the workplace. The OSHA website contains Fact Sheets and Guidance Documents that relate to heat and cold stress that have been incorporated into this program.

## 2.0 Responsibilities

### 2.1 EHS shall:

- 2.1.1 Maintain, review and update the Heat and Cold Stress Program as needed.
- 2.1.2 Provide monitoring (upon request) and assist employees with the development of procedures to minimize the adverse effects of heat and cold stress in the workplace.
- 2.1.3 Provide training to employees affected by heat and cold.

### 2.2 Supervisors shall:

- 2.2.1 Review and comply with the provisions outlined in this program.
- 2.2.2 Ensure all employees are properly trained before working in extreme temperature conditions.
- 2.2.3 Assess the day-to-day heat or cold stresses on employees.
- 2.2.4 Assess employees work load and assigning work and rest schedules as needed.
- 2.2.5 Ensure all employees have the appropriate personal protective equipment (PPE) prior to working in extreme temperature conditions.
- 2.2.6 Ensure employees are familiar with this safety program.

### 2.3 Employees shall:

- 2.3.1 Review and comply with the provisions outlined in this program.
- 2.3.2 Complete training before working in extreme temperature conditions.

- 2.3.3 Wear the appropriate PPE.
- 2.3.4 Report heat and cold stress concerns to their supervisor.

### **3.0 Heat Related Illnesses; Signs, Treatment and Prevention**

3.1 While working in hot weather conditions, the human body may not be able to maintain a normal temperature just by sweating. If this happens, heat-related illnesses may occur. The most common health problems caused by hot work environments include:

- 3.1.1 Heat stroke – This is the most serious heat related effect. Heat stroke occurs when the body temperature increases above 104<sup>0</sup>F. Signs and symptoms of heat stroke are confusion, loss of consciousness and lack of perspiration. This condition must be treated as a medical emergency and the employee must receive immediate medical attention.
- 3.1.2 Heat exhaustion – Signs and symptoms of heat exhaustion include headache, nausea, dizziness, weakness, irritability, confusion, thirst, heavy perspiration and a body temperature greater than 100.4<sup>0</sup>F. Employees experiencing heat exhaustion should be moved to a cool area, given fluids to drink and given cold compresses for their head, face and neck. Employees should also be taken to a clinic or emergency room to be monitored by medical personnel.
- 3.1.3 Heat cramps – Signs and symptoms of heat cramps include muscle pains usually caused by the loss of body salts/fluids. Employees should replace fluid loss by drinking water and/or carbohydrate-electrolyte replacement liquids (e.g. Gatorade) every 15 to 20 minutes.
- 3.1.4 Heat rash – Heat rash is caused by excessive perspiration and looks like a red cluster of pimples or small blisters. Heat rash usually appears on the neck, upper chest, in the groin, under the breasts and in elbow creases. Treatment for heat rash is to provide a cooler, less humid environment.
- 3.1.5 Dehydration – Dehydration is a major factor in most heat disorders. Signs and symptoms of dehydration include increasing thirst, dry mouth, weakness or light-headedness (particularly if worse upon standing), and a darkening of the urine or a decrease in urination. Dehydration can be reversed or put back in balance by drinking fluids that contain electrolytes (i.e. Gatorade) that are lost during work related activities. Avoid caffeinated drinks.

3.2 While heat related illness are dangerous and potentially life threatening, they can be prevented. Prevention methods include:

- 3.2.1 Acclimation – Acclimation is a process by which the physical processes of an employee's body adjusts to the environment over a period of time. Based on data obtained from OSHA, this process usually takes five to seven days. This process could take up to three weeks depending on the individual and their work environment. According to the American Industrial Hygiene Association, the process requires a consistent work level for at least two hours each day during the acclimation period in order for an employee to become acclimatized. Mere exposure to heat does not confer acclimatization, nor does acclimatization at one heat stress level confer resistance to heat stress at a higher temperature or more vigorous work load.

Employees who are not adequately acclimatized to the heat may experience temporary heat fatigue resulting in a decline in performance, coordination or alertness. They may also become irritable or depressed. This can be prevented through gradual adjustment to the hot

environment. People in good physical condition tend to acclimatize better because their cardiovascular systems respond better.

- 3.2.2 Engineering Controls – For employees working indoors, the best way to prevent heat-related illness is to make the work environment cooler. Where and if possible, use air conditioning to cool the work area. Alternatively, increase the general ventilation as much as possible by opening windows or doors. When available, use cooling fans to aid in increasing ventilation.
- 3.2.3 Safe Work Practices – For employees working outdoors or working indoors without air conditioning or ventilation, take scheduled breaks in cool areas. Ensure there is plenty of cool water to drink and take water breaks as needed. Immediately report any problems to a supervisor. Supervisors should consider scheduling the hottest work for the coolest part of day, assigning extra employees to high demand tasks, and using work-saving devices (e.g. power tools, hoists or lifting aids) to reduce the body's work load. All employees should watch out for the safety of their coworkers.
- 3.2.4 Heat Index – The Heat Index is a single numeric value that uses both temperature and humidity to inform the public on how the weather outdoors "feels". The higher the Heat Index, the hotter the weather feels. OSHA has used the Heat Index to assign protective measures for workers as the Heat Index increases. These protective measures may reduce the likelihood of heat related illnesses. The Heat Index and related protective measures are contained in Appendix A.

### 3.3 Wet Bulb Globe Temperature (WBGT) Monitoring.

- 3.3.1 Upon request, EHS can provide WBGT monitoring using a QuesTemp 32 Heat Stress Monitor. With this instrument, EHS personnel can provide supervisors and employees with the following information:
  - 3.3.1.1 Natural Wet Bulb Thermometer – Gives an indication of the effects of humidity on an individual.
  - 3.3.1.2 Globe Thermometer – Gives an indication of the radiant heat exposure on an individual due to either direct light or hot objects in the environment.
  - 3.3.1.3 Dry Bulb Thermometer – A measurement of the ambient air temperature.
  - 3.3.1.4 Relative Humidity – A measurement of water vapor in the air.
- 3.3.2 After collecting the aforementioned information, EHS can compare the results to published charts that show the allowable work-rest regimens for given workloads.

## 4.0 Cold Related Illnesses and Injuries; Signs, Treatment and Prevention

- 4.1 During cold weather, an employee's body will use energy to maintain a normal internal body temperature. This will result in a shift of blood flow from employee's extremities (hands, feet and legs) and outer skin to the employee's core (chest and abdomen). If this happens, cold-related illnesses and injuries may occur if exposed to cold conditions for an extended period of time. The most common health problems caused by cold work environments include:
  - 4.1.1 Hypothermia – Hypothermia is a potentially serious health condition. Hypothermia occurs when body heat is lost faster than it can be replaced. When the core body temperature drops

to approximately 95°F, the onset of symptoms normally begins. The employee may begin to shiver, lose coordination, have slurred speech, and fumble with items in the hand. The employee's skin will likely be pale and cold. As the body temperature continues to fall these symptoms will worsen and shivering will stop. Once the body temperature falls to around 85°F severe hypothermia will develop and the person may become unconscious, and at 78°F, vital organs may begin to fail.

Treatment depends on the severity of the hypothermia. For cases of mild hypothermia move to warm area and stay active. Remove wet clothes and replace with dry clothes or blankets, cover the head. To promote metabolism and assist in raising internal core temperature drink a warm (not hot) sugary drink. Avoid drinks with caffeine. For more severe cases do all the above, plus contact emergency medical personnel (Call 911 for an ambulance), cover all extremities completely, place very warm objects, such as hot packs or water bottles on the victim's head, neck, chest and groin. Arms and legs should be warmed last. In cases of severe hypothermia, treat the employee very gently and do not apply external heat to re-warm. Hospital treatment is required.

- 4.1.2 Frostbite – Frostbite occurs when the skin actually freezes and loses water. In severe cases, amputation of the frostbitten area may be required. While frostbite usually occurs when the temperatures are 30° F or lower, wind chill factors can allow frostbite to occur in above freezing temperatures. Frostbite typically affects the extremities, particularly the feet and hands. The affected body part will be cold, tingling, stinging or aching followed by numbness. Skin color turns red, then purple, then white, and is cold to the touch. There may be blisters in severe cases.

Do not rub the area to warm it. Wrap the area in a soft cloth, move the employee to a warm area, and contact medical personnel. Do not leave the employee alone. If help is delayed, immerse in warm (maximum 105 °F), not hot, water. Do not pour water directly on affected part. If there is a chance that the affected part will get cold again do not warm. Repeated heating and cooling of the skin may cause severe tissue damage.

- 4.1.3 Trench Foot – Trench Foot is caused by having feet exposed to damp, unsanitary and cold conditions including water at temperatures above freezing for long periods of time. It is similar to frostbite, but considered less severe. Symptoms usually consist of tingling, itching or burning sensation. Blisters may be present.

For treatment, soak feet in warm water, then wrap with dry cloth bandages. Drink a warm, sugary drink. Seek medical attention if necessary.

- 4.1.4 Dehydration – It is easy to become dehydrated during cold weather. Signs of dehydration include increasing thirst, dry mouth, weakness or light-headedness (particularly if worse upon standing), and a darkening of the urine or a decrease in urination. Dehydration can be reversed or put back in balance by drinking fluids that contain electrolytes (i.e. Gatorade) that are lost during work related activities. Avoid caffeinated drinks

- 4.2 Just as with heat related illness, cold related illnesses and injuries are dangerous and potentially life threatening, however, they can be prevented. Prevention methods include:

- 4.2.1 Acclimation – Employees exposed to the cold should be physically fit, without any circulatory, metabolic, or neurologic diseases that may place them at increased risk for hypothermia. A new employee should not be required to work in the cold full time during the first days of employment until they become adjusted to the working conditions and required protective

clothing. New employees should be introduced to the work schedule slowly and be trained accordingly.

- 4.2.2 Engineering Controls – For employees working indoors, the best way to prevent cold-related illness is to make the work environment warmer. Where and if possible, use heaters to warm the work area. Alternatively, decrease the general ventilation as much as possible by closing windows or doors.
- 4.2.3 Safe Work Practices – For employees working outdoors or working indoors without heat, take scheduled breaks in warm areas. If available, use wind barricades to block the wind from the employees. Ensure there is plenty of water to drink and take water breaks as needed. Immediately report any problems to a supervisor. Supervisors should consider scheduling the most work for the warmest part of day, assigning extra employees to high demand tasks that will require longer periods in cold areas. All employees should watch out for the safety of their coworkers.
- 4.2.4 Personal Protective Equipment (PPE) – PPE is an important factor in preventing cold stress related illnesses and injuries. Employees should adhere to the following recommendations when dressing for work in a cold environment:
- Wear at least three layers of clothing; an inner layer of wool, silk or synthetic to wick moisture away from the body; a middle layer of wool or synthetic to provide insulation even when wet; an outer wind and rain protection layer that allows some ventilation to prevent overheating.
  - Wear a hat or hood; up to 40% of body heat can be lost when the head is left exposed.
  - Wear insulated boots or other footwear.
  - Do not wear tight clothing; loose clothing provides better ventilation.
  - Keep a change of clothing available in case work clothes become wet.
- 4.2.5 The Cold Stress Equation – OSHA has incorporated information obtained from the American Conference of Governmental Industrial Hygienists (ACGIH) threshold limit values into the Cold Stress Equation. As the temperature decreases and/or the wind speed increases, the potential for cold stress related illnesses and injuries increases. The Cold Stress Equation is contained in Appendix B.

## 5.0 Training

- 5.1 Supervisors shall ensure all employees have received Heat and/or Cold Stress training prior to working in such conditions.
- 5.2 EHS can provide heat or cold stress training upon request.

## 6.0 Recordkeeping

- 6.1 All training records should be maintained in the employees personnel file and maintained by the supervisor. Training records are maintained in EH&S for training programs provided by EH&S.

## **Appendix A**

### **Heat Index**

The heat index is a simple tool and a useful guide for employers/employees making decisions about protecting employees in hot weather. It does not account for certain conditions that contribute additional risk, such as physical exertion. Consider taking the steps at the next highest risk level to protect employees from the added risks posed by:

- Working in the direct sun (can add up to 15°F to the heat index value)
- Wearing heavy clothing or protective gear

Under most circumstances, fluid intake should not exceed 6 cups per hour or 12 quarts per day. This makes it particularly important to reduce work rates, reschedule work, or enforce work/rest schedules.

The heat index follows on the next two pages.



Heat Index	Risk Level	Protective Measures
<91°F	Lower  (Caution)	<ul style="list-style-type: none"> <li>• Provide plenty of drinking water</li> <li>• Ensure that adequate medical services are available</li> <li>• Plan ahead for times when heat index is higher, including worker heat safety training</li> <li>• Encourage workers to wear sunscreen</li> <li>• If workers must wear heavy protective clothing , perform strenuous activity or work in the direct sun, addition precautions are recommended to protect workers from heat related illness</li> </ul>
91°F to 103°F	Moderate	In addition to the steps listed above: <ul style="list-style-type: none"> <li>• Remind workers to drink water often (about 4 cups per hour)</li> <li>• Review heat related illness topics with workers such as recognition, prevention and first-aid</li> <li>• Schedule frequent breaks in cool, shaded areas</li> <li>• Acclimatize workers</li> <li>• Set up a buddy system and instruct workers and supervisors to watch for signs of heat related illnesses</li> <li>• Schedule strenuous activities at a time when the heat index is lower</li> <li>• Develop work rest schedules</li> <li>• Monitor workers closely</li> <li>• If workers must wear heavy protective clothing , perform strenuous activity or work in the direct sun, addition precautions are recommended to protect workers from heat related illness</li> </ul>

Heat Index	Risk Level	Protective Measures
103°F to 115°F	High	<p>In addition to the steps listed above:</p> <ul style="list-style-type: none"> <li>• Alert workers of high risk conditions</li> <li>• Actively encourage workers to drink plenty of water (about 4 cups per hour)</li> <li>• Limit physical exertion</li> <li>• have a knowledgeable person at the work site who is well informed about heat related illness and able to determine appropriate work/rest schedules</li> <li>• Establish and enforce work/rest schedules</li> <li>• Adjust work activities (e.g. reschedule work, pace/rotate jobs)</li> <li>• Use cooling techniques</li> <li>• Watch/communicate with workers at all times</li> <li>• When possible, reschedule activities to a time when the heat index is lower</li> </ul>
115°F	Very High to Extreme	<p>If essential work must be done, in addition to the steps listed above:</p> <ul style="list-style-type: none"> <li>• Alert workers of extreme heat hazards</li> <li>• Establish water drinking schedule (about 4 cups per hour)</li> <li>• Develop and enforce protective work/rest schedules</li> <li>• Conduct physiological monitoring (e.g. pulse, temperature, etc.)</li> <li>• Stop work if essential control methods are inadequate or unavailable</li> <li>• Reschedule non-essential activities for days with a reduced heat index or to a time when the heat index is lower</li> <li>• Move essential work tasks to the coolest part of the work shift</li> <li>• Consider earlier start times, split shifts or evening/night shifts</li> <li>• Strenuous work tasks and those requiring the use of heavy or non-breathable clothing or impermeable chemical protective clothing should not be conducted when the heat index is at or above 115°F</li> </ul>

Appendix B

The Cold Stress Equation

# THE COLD STRESS EQUATION

LOW TEMPERATURE + WIND SPEED + WETNESS  
= INJURIES & ILLNESS

